

## EX-VESSEL VALUE AND SIZE COMPOSITION OF REPORTED MAY-AUGUST CATCHES OF BROWN SHRIMP AND WHITE SHRIMP FROM 1960 TO 1981 AS RELATED TO THE TEXAS CLOSURE<sup>1</sup>

CHARLES WAX CAILLOUET, Jr. AND DENNIS BRIAN KOI  
National Marine Fisheries Service, Southeast Fisheries Center,  
Galveston Laboratory, Galveston, Texas 77550

**ABSTRACT** Indices were used to test for trends in ex-vessel price spread (value per shrimp by size category), size composition, and ex-vessel value composition of the reported May-August catches (inshore and offshore combined) of brown shrimp (*Penaeus aztecus*) and white shrimp (*P. setiferus*) from the Texas coast, the Mississippi River to Texas, and Pensacola to the Mississippi River, from 1960 to 1981. Levels of reported May-August catch and ex-vessel value of the catch also were examined for the same period. Statistical tests were conducted to determine if 1981 was an outlier as compared to other years, in the context of impacts of closure of the fishery conservation zone (FCZ) off Texas to shrimping from May 22 to July 15, 1981, a management measure referred to as the Texas Closure.

### INTRODUCTION

The territorial sea of the State of Texas and the adjacent fishery conservation zone (FCZ) were closed to shrimping for brown shrimp (*Penaeus aztecus*) from May 22 to July 15, 1981. This management action was dubbed the Texas Closure by the Gulf of Mexico Fishery Management Council (GMFMC 1980). The territorial sea extends from the coastal baseline to 9 nautical miles off Texas (Figure 1). The FCZ, under federal jurisdiction, extends from the outer limit of Texas' territorial sea to 200 miles from shore. The State of Texas eliminated minimum size restrictions on brown shrimp caught in waters open to shrimping before and after the closure, and allowed daytime fishing for white shrimp to continue in waters up to 4 fm in depth within the territorial sea during the closure. The GMFMC (1980) expected that the Texas Closure would lead to an increase in yield of brown shrimp in the FCZ through additional growth and survival and from abatement of the practice of discarding undersized brown shrimp.

For years, there has been considerable controversy regarding the optimum size at which shrimp should be harvested (GMFMC 1980). During their life cycle, brown shrimp and white shrimp enter inshore waters (landward of barrier islands) as postlarvae, and emigrate a few months later to offshore waters (seaward of barrier islands) as subadults. For that reason, the shrimp caught inshore generally are smaller than those caught offshore. Thus, to a great extent, the size composition of the catch is dependent on where and when the shrimp are caught during each annual cycle. This choice is complicated by the fact that ecological requirements of brown shrimp and white shrimp differ from one another (Christmas and Etzold 1977, GMFMC 1980), and the peaks in abundance of the two species occur out of phase with one




Figure 1. Boundaries of statistical areas 10-21, the Texas territorial sea and the fishery conservation zone off Texas (based on information from GMFMC, 1980).

another. The size at which shrimp are harvested is all the more important, because the ex-vessel price per pound of shrimp increases with their size (Neal 1967, Griffin et al. 1974, Griffin and Nichols 1976, Griffin et al. 1976).

The 1981 Texas Closure provided a unique opportunity for exaggerating the contrast between a management strategy that protects small brown shrimp and allows them to grow to larger sizes before harvest and one that allows small as well as large brown shrimp to be harvested in large quantities. Using yield-per-recruit analysis and a simulation model of shrimp fishing, Nichols (1982) predicted that the brown shrimp yield off Texas in May-August 1981 would be 11.7 million pounds greater than that expected had there been no Texas Closure. Poffenberger (1982) showed that the Texas Closure caused an increase in brown shrimp landings off the Texas coast, which in turn caused a decrease in ex-vessel price per pound. Despite the decrease in price, Poffenberger (1982) concluded that there was an increase in gross revenue by about 21.5 million dollars from the brown

<sup>1</sup>Contribution No. 82-34G from the Southeast Fisheries Center, Galveston Laboratory, National Marine Fisheries Service, NOAA.

Manuscript received Sept. 27, 1982; accepted Dec. 3, 1982.

shrimp fishery in May-August 1981 over what would have been expected had there been no Texas Closure. Klima et al. (1982) stated that recruitment of brown shrimp in Texas bays and size composition of the offshore stock in 1981 were similar to those in previous years of good production. They also stated that the brown shrimp catch per unit effort was higher in 1981 than would have been the case without the Texas Closure. They concluded that the Texas Closure had a positive impact on relative abundance and production of brown shrimp off Texas. Using fishery-independent data, Matthews (1982) reported good catch rates of brown shrimp collected off the coast of Texas during the Texas Closure.

Caillouet and Koi (1981) investigated trends in ex-vessel price spread among shrimp size categories, in size composition, and in ex-vessel value composition of the May-August catches (inshore and offshore combined) of brown shrimp and white shrimp from the Texas, Louisiana, Mississippi, and Alabama coasts, for the years 1960 to 1978. They anticipated use of their analytical approach in assessing future impacts of the Texas Closure. When they chose May-August as the time interval for their analysis, the specific dates for the Texas Closure had not yet been set. They wanted to make sure that the chosen interval would encompass the future closure interval so as to assess immediate impacts of the Texas Closure, including those immediately before and in anticipation of the closure, and those during and immediately after the closure.

While almost half of the May-August period was closed to shrimping for brown shrimp off the Texas coast in 1981, it was open to shrimping in other areas, both inshore and offshore. Caillouet and Koi (1981) based their analyses on inshore and offshore catch data combined. The stocks of shrimp include inshore and offshore components. The possibility that inshore fishing intensity might increase in areas open to shrimping during the Texas Closure (GMFMC 1980), with consequent effects on the combined inshore and offshore catch, provided further justification for combining inshore and offshore catches in the analysis. Furthermore, because the yield from inshore areas represents a significant portion of the total yield, we thought that it should not be ignored in assessing impacts of the Texas Closure. In other words, we did not want to assume that impacts of the Texas Closure were confined only to the offshore fishery.

Two purposes of this paper are to update the analyses of Caillouet and Koi (1981) by adding data from 1979 to 1981, and to compare the observed 1981 indices of ex-vessel price spread, size composition, and ex-vessel value composition of the May-August catches of brown shrimp and white shrimp with those expected for 1981 based on average levels or trends from 1960 to 1981. For the latter purpose, we applied linear trend analyses, employing simple linear regression, to the time series of indices over the years 1960–1981. Our analyses were designed to test the null hypothesis that

the 1981 indices were not significant<sup>2</sup> “outliers” in the 1960–1981 data series.

Our application of simple linear regression analysis, to detect trends and to test hypotheses concerning indices of ex-vessel price spread, size composition, and ex-vessel value composition of shrimp catches, requires some explanation to guide the reader in interpretation of the results. Simple linear regression analysis tests only for linear trends. If there were strongly curvilinear trends in the indices, linear regression analysis alone would not detect them, and this could also lead to a conclusion that there were no significant (linear) trends. However, inspection of our scatter plots (e.g., Figure 9) of the data points by the reader can be helpful in this regard. When our analysis detected no significant linear trend, a mean index was calculated to represent the time series of points.

As shown in our scatter plots of the indices, two types of 95 percent confidence bands were given. The narrower confidence bands (Figure 9) represented the uncertainty in estimating or predicting the population mean index, depicted by either a regression line or a horizontal line through the data points (for cases where there was no significant linear trend; e.g., Figure 10). The wider confidence bands (Figure 9) represented the uncertainty in estimating or predicting an individual index level for a particular year. In general, 5 percent of the points would be expected to fall outside such confidence bands, due to chance alone. For additional explanation, the reader is referred to Snedecor and Cochran (1967, pp. 153–157).

A third purpose of this paper is to determine whether the three indices were correlated with each other or with the weight of the May-August catches, using the entire time series of indices from 1960 to 1981. We made comparisons among such correlations between species and among coastal areas, in order to distinguish the effects of the Texas Closure from other effects such as year-to-year variations in recruitment, fishing effort, or both.

## METHODS

### *Description of data*

Summations of the May-August catches (inshore and offshore combined) for brown shrimp and white shrimp and their ex-vessel value were compiled from data files available from the National Marine Fisheries Service (NMFS), Southeast Fisheries Center (SEFC), Technical and Information Management Services (TIMS). The data for 1979, 1980, and 1981 were those available from the TIMS in February 1982. The reader is cautioned that the 1980 and 1981 data may have undergone slight changes by the TIMS since then, based upon identification of minor errors and upon minor additions

<sup>2</sup> Refers throughout this paper to the 95% level of confidence.

to the data files. Historically, such updating of the files has not been of major consequence.

The characteristics of shrimp catch data were described previously (Caillouet and Koi 1981). Data for 1960 to 1978 were obtained from Caillouet and Koi (1981). The weight of the reported May-August catches (inshore and offshore combined) was expressed in pounds (heads off) and the ex-vessel value in dollars (unadjusted for effects of inflation) for each year, coastal area, species, and size category (< 15, 15–20, 21–25, 26–30, 31–40, 41–50, 51–67, and  $\geq$  68 count), and "pieces," representing parts of shrimp tails that could not be assigned to a size category.

Separate analyses were conducted for both species and three coastal areas (Figure 1), which were distinguished as follows: 1) Texas coast (statistical areas 18–21 combined); 2) Mississippi River to Texas (statistical areas 13–17 combined), representing that part of the Louisiana coast west of the Mississippi River; and 3) Pensacola to the Mississippi River (statistical areas 10–12 combined), representing that part of the Louisiana coast east of the Mississippi River, the Mississippi coast, the Alabama coast, and a small part of the upper west coast of Florida (catches from Pensacola Bay are excluded from this area, as they are allocated to the adjacent Apalachicola area by the TIMS). Note that almost half of statistical area 17 was included in the area that was closed to shrimping from May 22 to July 15, 1981 (Figure 1). Therefore, for the years 1960 to 1980, the May-August catch statistics for the Mississippi River to Texas coastal area represent a somewhat larger zone open to shrimping than was the case in 1981, as a result of the Texas Closure.

#### ANALYTICAL RESULTS

The analytical methods followed those of Caillouet and Koi (1981).

##### May-August catches

Table 1 provides background concerning both the magnitude and stability of the percentage of the May-August catch (all size categories as well as "pieces" combined) taken inshore vs. offshore, for each species and coastal area from 1960 to 1981. For brown shrimp, the percentage of the May-August catch taken inshore was lowest for the Texas coast (10%), intermediate for Pensacola to the Mississippi River (50%), and highest for the Mississippi River to Texas (54%). This elucidates the emphasis on offshore fishing in Texas and on both inshore and offshore fishing in the other two coastal areas. However, the inshore proportion of the May-August brown shrimp catch has been increasing on the Texas coast and in the Pensacola to the Mississippi River area, while it has been decreasing in the Mississippi River to Texas area. For white shrimp, the percentage of the May-August catch taken inshore was highest for Pensacola to the Mississippi River (62%), intermediate for the Texas coast (54%), and lowest for the Mississippi River to Texas (32%). This percentage showed no significant trend for the Texas coast and Mississippi River to Texas, but there was a significant downward

TABLE 1.

Percentages of the May-August catches (inshore and offshore combined) of brown shrimp and white shrimp taken inshore, and percentages of the calendar year annual catches (inshore and offshore combined) of brown shrimp and white shrimp taken during May-August, for the Texas Coast, the Mississippi River to Texas, and Pensacola to the Mississippi River, during 1960 to 1981.

Species	Coastal Area	Percentage May-August Catch Taken Inshore		Percentage Annual Catch Taken in May-August	
		Mean <sup>a</sup>	Range	Mean <sup>a</sup>	Range
Brown Shrimp	Texas Coast	10 <sup>b</sup>	0 <sup>c</sup> –29	58	47–73
	Mississippi River to Texas	54 <sup>d</sup>	35–68	81	74–90
	Pensacola to Mississippi River	50 <sup>e</sup>	34–76	84	75–93
White Shrimp	Texas Coast	54	30–78	24	16–34
	Mississippi River to Texas	32	11–47	19	8–25
	Pensacola to Mississippi River	62 <sup>f</sup>	29–94	13	5–23

<sup>a</sup> Averaged over all years from 1960 to 1981.

<sup>b</sup> There was a significant upward trend in percentage from 1960 to 1981:  $Y = -46.2 + 0.8X$ , where  $Y$  = percentage, and  $X$  = last two digits (60–81) of each year (1960–1981). The coefficient of determination,  $r^2$ , was 0.55.

<sup>c</sup> Rounded to zero, but the percentage was 0.04.

<sup>d</sup> There was a significant downward trend in percentage from 1960 to 1981:  $Y = 109.0 - 0.8X$ . See footnote b for definitions of  $Y$  and  $X$ . The coefficient of determination,  $r^2$ , was 0.26.

<sup>e</sup> There was a significant downward trend in percentage from 1960 to 1981:  $Y = -41.6 + 1.3X$ . See footnote b for definitions of  $Y$  and  $X$ . The coefficient of determination,  $r^2$ , was 0.61.

<sup>f</sup> There was a significant downward trend in percentage from 1960 to 1981:  $Y = 206.2 - 2.0X$ . See footnote b for definitions of  $Y$  and  $X$ . The coefficient of determination,  $r^2$ , was 0.50.

trend in this percentage for Pensacola to the Mississippi River.

The percentage of the calendar year annual catch of brown shrimp represented by the May-August catch of brown shrimp is shown in Table 1, for each coastal area. Likewise, a similarly calculated percentage is shown for white shrimp. There were no significant trends in this percentage for either species in any of the three coastal areas from 1960 to 1981. For Pensacola to the Mississippi River, the May-August brown shrimp catch represented 84% of the annual catch, on the average. For the Mississippi River to Texas it averaged 81%, and for the Texas coast 58%. Thus, a substantial percentage of the annual brown shrimp catch was taken in May-August within each coastal area. The lower percentage for the Texas coast reflects the emphasis on offshore fishing with its consequent protracted harvest. For

white shrimp, the average percentage of the annual catch taken in May-August was highest for the Texas coast (24%), intermediate for the Mississippi River to Texas (19%), and lowest for Pensacola to the Mississippi River (13%). These low percentages reflected the later harvest of white shrimp, compared to brown shrimp, determined by differences in phasing of life cycle events in these two species.

Within each coastal area, the May-August catch of brown shrimp greatly exceeded that of white shrimp in all years from 1960 to 1981 (Figures 2-7). In all years, the May-August brown shrimp catches from the Mississippi River to




Figure 2. Weight (millions of pounds, heads off) and ex-vessel value (millions of dollars) of reported May-August catches (inshore and offshore combined) of brown shrimp from the Texas coast (statistical areas 18-21 combined), 1960-1981.

Texas exceeded those from Pensacola to the Mississippi River, and in most years, including 1981, they exceeded those from the Texas coast. The May-August catch of brown shrimp from the Texas coast was higher in 1981 than in any previous year except 1967, and the ex-vessel value of this catch was at an unprecedented high for the Texas coast (Figure 2). The May-August brown shrimp catches from the other two coastal areas in 1981 were higher than in 1980, but did not exceed those of all previous years in the 1960-1980 series (Figures 4 and 6).

Despite the fact that the May-August brown shrimp catch from the Texas coast (29.6 million lb., Figure 2) in 1981 was lower than that from the Mississippi River to Texas (38.4 million lb., Figure 4), it had a greater ex-vessel value (66.3 million dollars) than that from the Mississippi River to Texas (56.8 million dollars). In 1981, the ex-vessel value (19.3 million dollars) of the May-August brown shrimp catch from Pensacola to the Mississippi River (10.4 million lb.) was much lower than for the other two areas, as was the case in all previous years.

The May-August catch of white shrimp from the Texas coast (1.7 million lb., Figure 3) in 1981 was lower than that




Figure 3. Weight (millions of pounds, heads off) and ex-vessel value (millions of dollars) of reported May-August catches (inshore and offshore combined) of white shrimp from the Texas coast (statistical areas 18-21 combined), 1960-1981.




Figure 4. Weight (millions of pounds, heads off) and ex-vessel value (millions of dollars) of reported May-August catches (inshore and offshore combined) of brown shrimp from the Mississippi River to Texas (statistical areas 13-17 combined), 1960-1981.

from the Mississippi River to Texas (7.8 million lb., Figure 5), and the ex-vessel value of this catch (5.6 million dollars) was dramatically lower than that from the Mississippi River to Texas (26.6 million dollars). The May-August white shrimp catches from the Mississippi River to Texas and from Pensacola to the Mississippi River in 1981 were somewhat higher than in 1980, but the opposite was true for the Texas coast. In all years, the May-August white shrimp catch from the Mississippi River to Texas exceeded that from the Texas coast, and both exceeded that from Pensacola to the Mississippi River. The ex-vessel value of the May-August catch of white shrimp from the Mississippi River to Texas in 1981 was at an all time high. While the ex-vessel value of the






Figure 5. Weight (millions of pounds, heads off) and ex-vessel value (millions of dollars) of reported May-August catches (inshore and offshore combined) of white shrimp from the Mississippi River to Texas (statistical areas 13-17 combined), 1960-1981.




Figure 6. Weight (millions of pounds, heads off) and ex-vessel value (millions of dollars) of reported May-August catches (inshore and offshore combined) of brown shrimp from Pensacola to the Mississippi River (statistical areas 10-12 combined), 1960-1981.

May-August catch of white shrimp from Pensacola to the Mississippi River in 1981 was higher than that in most years in the 1960-1980 series, it did not exceed that in 1977 and 1979.

*May-August ex-vessel value per shrimp by size category*

Caillouet and Koi (1981) used the logarithmic form of the following model to estimate parameters a and b by linear regression, for each species, coastal area, and year from 1960 to 1978, and we updated this series through 1981 (Tables 2-4):

$$\hat{V}_i = a (\exp bC_i) \quad (1)$$

where  $V_i$  = May-August average ex-vessel value (dollars per




Figure 7. Weight (millions of pounds, heads off) and ex-vessel value (millions of dollars) of reported May-August catches (inshore and offshore combined) of white shrimp from Pensacola to the Mississippi River (statistical areas 10-12 combined), 1960-1981.

shrimp) for the  $i$ th size category,  $C_i$  = lower limit (count) of the  $i$ th size category ( $C_1 = 15, C_2 = 21, C_3 = 26, C_4 = 31, C_5 = 41, C_6 = 51, \text{ and } C_7 = 68$ ), and  $i = 1, 2, \dots, 7$ . An example of the relationship between  $V$  and  $C$  is shown in Figure 8A, based on the May-August brown shrimp catch from the Texas coast in 1981. An example of the fitted logarithmic form of model 1 is given in Figure 8B. The very high coefficients of determination,  $r^2$ , for the logarithmic form of model 1 indicated that straight lines fitted the data for each year very well (Tables 2-4, see also Caillouet and Koi 1981). All slopes,  $b$ , of the straight lines were negative, showing that the ex-vessel price per shrimp decreased logarithmically with increase in count (decrease in size). The slope,  $b$ , of each straight line is a simple index of the May-August ex-vessel price spread among the size categories of shrimp in any given year.

TABLE 2.

Parameter estimates<sup>a</sup> of linear regressions of transformed ex-vessel price (dollars per shrimp),  $\ln V$ , on count,  $C$ , for reported May-August catches (inshore and offshore combined) of brown shrimp and white shrimp from the Texas coast (statistical areas 18-21 combined), 1979-1981.

Year	Brown Shrimp <sup>b</sup>			White Shrimp <sup>b</sup>		
	a	b	r <sup>2</sup>	a	b	r <sup>2</sup>
1979	0.80614	-0.05504	0.996	0.70034	-0.05206	0.979
1980	0.59210	-0.05101	0.994	0.49137	-0.04706	0.985
1981	0.51354	-0.05318	0.966	0.60401	-0.05432	0.948

<sup>a</sup>  $\ln(a)$  = intercept,  $b$  = slope, and  $r^2$  = coefficient of determination.  
<sup>b</sup> Results of similar analyses of data from 1960 to 1978 can be found in Caillouet and Koi (1981).

TABLE 3.

Parameter estimates<sup>a</sup> of linear regressions of transformed ex-vessel price (dollars per shrimp),  $\ln V$ , on count,  $C$ , for reported May-August catches (inshore and offshore combined) of brown shrimp and white shrimp from the Mississippi River to Texas (statistical areas 13–17 combined), 1979–1981.

Year	Brown Shrimp <sup>b</sup>			White Shrimp <sup>b</sup>		
	a	b	r <sup>2</sup>	a	b	r <sup>2</sup>
1979	0.77312	-0.05398	0.995	0.81161	-0.05504	0.990
1980	0.56006	-0.05004	0.995	0.56161	-0.05018	0.992
1981	0.57987	-0.05529	0.976	0.73555	-0.05975	0.976

<sup>a</sup>  $\ln(a)$  = intercept,  $b$  = slope, and  $r^2$  = coefficient of determination.

<sup>b</sup> Results of similar analyses of data from 1960 to 1978 can be found in Caillouet and Koi (1981).

TABLE 4.

Parameter estimates<sup>a</sup> of linear regressions of transformed ex-vessel value (dollars per shrimp),  $\ln V$ , on count,  $C$ , for reported May-August catches (inshore and offshore combined) of brown shrimp and white shrimp from Pensacola to the Mississippi River (statistical areas 10–12 combined), 1979–1981.

Year	Brown Shrimp <sup>b</sup>			White Shrimp <sup>b</sup>		
	a	b	r <sup>2</sup>	a	b	r <sup>2</sup>
1979	0.73353	-0.05047	0.989	0.73061	-0.05045	0.980
1980	0.53505	-0.04767	0.987	0.52388	-0.04739	0.985
1981	0.56535	-0.05258	0.972	0.62225	-0.05482	0.944

<sup>a</sup>  $\ln(a)$  = intercept,  $b$  = slope, and  $r^2$  = coefficient of determination.

<sup>b</sup> Results of similar analyses of data from 1960 to 1978 can be found in Caillouet and Koi (1981).

We excluded the < 15 size category from model 1 so as to be consistent with previous work, and because the line representing the logarithmic form of model 1 is not straight in the region of < 15 count (Caillouet and Koi 1981). The < 15 category represented  $\leq$  3% of the weight of the May-August catches of brown shrimp in any of the three coastal areas in any given year. In 1981, the < 15 category represented 0.2% of the May-August catches of brown shrimp from the Texas coast, 0.2% from the Mississippi River to Texas, and 0.1% from Pensacola to the Mississippi River. In any given year, the < 15 category represented as high as 23% of the May-August white shrimp catches from the Texas coast (15.6% in 1981), 15% from the Mississippi River to Texas (5.1% in 1981), and 28% from Pensacola to the Mississippi River (3.6% in 1981). The category "pieces" also was excluded from model 1, because it represents parts of shrimp tails which could not be assigned to a size category.

To determine whether there were significant trends in the index of ex-vessel price spread among size categories, we calculated the linear regression of  $b$  on the last two digits

(60–81) of each year from 1960 to 1981, for both species and all three coastal areas (Table 5). There were significant downward trends in  $b$  for brown shrimp in all three coastal areas (Figures 9, 11, and 13) and for white shrimp from the Mississippi River to Texas (Figure 12) and Pensacola to the Mississippi River (Figure 14), but there was no significant trend in  $b$  for white shrimp from the Texas coast (Figure 10).

The downward trends indicated that the May-August ex-vessel price spread among size categories increased during the period from 1960 to 1981. Because there was no significant trend in  $b$  for white shrimp from the Texas coast, we calculated the mean of the indices for the years 1960–1981 and plotted the mean as a horizontal solid line through the points in Figure 10. No data point for 1972 was included in calculating the regression trend for white shrimp from Pensacola to the Mississippi River, because no catch was reported for the  $\geq$  68 count category in 1972 (Caillouet and Koi 1981).

Within each coastal area, the ex-vessel price spread index  $b$  for brown shrimp in May-August 1981 fell within the 95% confidence bands for individual levels of  $b$  (Figures 9, 11, and 13), indicating that there was no significant departure in 1981 from the expected increase in ex-vessel price spread




Figure 8. Relationship between the ex-vessel price (dollars per shrimp),  $V$ , and count,  $C$ , for the reported May-August catch (inshore and offshore combined) of brown shrimp from the Texas coast (statistical areas 18–21 combined) in 1982 (graph A). In graph B, the line was fitted by linear regression (see Table 2).

TABLE 5.

Results of tests for trends<sup>a</sup> in ex-vessel price spread index b, in size composition index d, and in ex-vessel value composition index h, for reported May-August catches (inshore and offshore combined) of brown shrimp and white shrimp from the Texas coast, the Mississippi River to Texas, and Pensacola to the Mississippi River, during 1960 to 1981 (indices b, d and h are defined in the text).

Species	Coastal Area		For Ex-vessel Price Spread Index b	For Size Composition Index d	For Ex-vessel Value Composition Index h
Brown Shrimp	Texas Coast	Trend <sup>a</sup>	-0.0003 <sup>b</sup>	0.0024 <sup>b</sup>	0.0023 <sup>b</sup>
		Intercept	-0.0332	-0.2183	-0.2353
		Trend Coefficient of Determination	0.222	0.641	0.624
		Sum of Squares <sup>c</sup>	0.000290	0.002776	0.002932
Brown Shrimp	Mississippi River to Texas	Trend	-0.0005 <sup>b</sup>	0.0003 <sup>b</sup>	0.0002
		Intercept	-0.0187	-0.0273	-0.0281
		Trend Coefficient of Determination	0.367	0.334	0.065
		Sum of Squares	0.000376	0.000120	0.000317
Brown Shrimp	Pensacola to Mississippi River	Trend	-0.0006 <sup>b</sup>	0.0008 <sup>b</sup>	0.0006 <sup>b</sup>
		Intercept	-0.0091	-0.0798	-0.0711
		Trend Coefficient of Determination	0.486	0.396	0.221
		Sum of Squares	0.000319	0.000904	0.001067
White Shrimp	Texas Coast	Trend	-0.0002	0.0007	0.0007
		Intercept	-0.0349	-0.1015	-0.1137
		Trend Coefficient of Determination	0.092	0.061	0.035
		Sum of Squares	0.000499	0.007437	0.011162
White Shrimp	Mississippi River to Texas	Trend	-0.0006 <sup>b</sup>	0.0009 <sup>b</sup>	0.0004
		Intercept	-0.0124	-0.1076	-0.0879
		Trend Coefficient of Determination	0.361	0.215	0.043
		Sum of Squares	0.000488	0.002834	0.003594
White Shrimp	Pensacola to Mississippi River	Trend	-0.0005 <sup>b</sup>	-0.0003	-0.0009
		Intercept	-0.0143	-0.0163	0.0144
		Trend Coefficient of Determination	0.296	0.025	0.168
		Sum of Squares	0.000514	0.002274	0.003297

<sup>a</sup> For each species and coastal area, the trends are represented by the slopes of the linear regressions of indices b, d, and h, respectively, on X, where X represents the last two digits (60–81) of each year, 1960–1981. The indices b, d, and h are shown in Tables 2–4, 6–8, and 9–11, respectively, for the years 1979–1981, and in Caillouet and Koi (1981) for the years 1960–1978. Data for 1972 were excluded from regressions for white shrimp from Pensacola to the Mississippi River (see Caillouet and Koi 1981).

<sup>b</sup> Trend (slope) was significantly different from zero at the 95% level of confidence.

<sup>c</sup> Sum of squares of deviations from linear regression.

for brown shrimp in any of the three coastal areas. The same was true for the May-August 1981 ex-vessel price spread index for white shrimp from the Mississippi River to Texas and from Pensacola to the Mississippi River (Figures 12 and 14). The May-August 1981 ex-vessel price spread index for white shrimp from the Texas coast fell within the 95% confidence limits for individual levels of b (Figure 10), indicating that there was no significant departure in 1981 from the expected price spread based on the mean price spread index. Thus, based upon the inherent variations and trends in ex-vessel price spread, the May-August price spread

index was not a significant outlier in the 1960–1981 time series.

#### May-August cumulative catch by size category

Caillouet and Koi (1981) used the logarithmic form of the following model to estimate parameters c and d by linear regression for each species, coastal area, and year from 1960 to 1978, and we updated this through 1981 (Tables 6–8):

$$\hat{P}_i = c(\exp dC_i) \quad (2)$$

where  $P_i$  = cumulative weight (pounds, heads off) of the




Figure 9. Trend (solid line) in the ex-vessel price spread index  $b$  (open circles) for reported May-August catches (inshore and offshore combined) of brown shrimp from the Texas coast (statistical areas 18–21 combined), 1960–1981 (data from Tables 2 and 5, and Caillouet and Koi 1981). The 95% confidence bands are shown for the trend line by short dashes and for individual values of  $b$  by long dashes.




Figure 11. Trend (solid line) in the ex-vessel price spread index  $b$  (open circles) for reported May-August catches (inshore and offshore combined) of brown shrimp from the Mississippi River to Texas (statistical areas 13–17 combined), 1960–1981 (data from Tables 3 and 5, and Caillouet and Koi 1981). The 95% confidence bands are shown for the trend line by short dashes and for individual values of  $b$  by long dashes.




Figure 10. Mean (solid line) ex-vessel price spread index  $b$  (open circles) for reported May-August catches (inshore and offshore combined) of white shrimp from the Texas coast (statistical areas 18–21 combined), 1960–1981 (data from Table 2, and Caillouet and Koi 1981). The 95% confidence limits are shown for the mean by short dashes and for individual values of  $b$  by long dashes.




Figure 12. Trend (solid line) in the ex-vessel price spread index  $b$  (open circles) for reported May-August catches (inshore and offshore combined) of white shrimp from the Mississippi River to Texas (statistical areas 13–17 combined), 1960–1981 (data from Tables 3 and 5, and Caillouet and Koi 1981). The 95% confidence bands are shown for the trend line by short dashes and for individual values of  $b$  by long dashes.

May-August catch in the  $i$ th size category. The catches in each size category were cumulated starting with the size category of smallest shrimp (highest count,  $\geq 68$ ) and continuing toward the size category of largest shrimp (lowest count, 15–20). An example of the relationship between  $P$  and  $C$  is shown in Figure 15A, for brown shrimp from the Texas coast in 1981. An example of the fitted logarithmic form of model 2 is shown in Figure 15B. The coefficients of determination for the straight lines representing each year

were very high (Tables 6–8, see also Caillouet and Koi 1981). All slopes,  $d$ , were negative, which reflected the construction of model 2 by cumulating catches from small-shrimp to large-shrimp size categories (Caillouet and Koi 1981). The slope,  $d$ , of each straight line is a simple index of the size composition of the May-August catch in any given year.

In each year, the levels of  $d$  for brown shrimp were lowest for the Texas coast, intermediate for Pensacola to





Figure 13. Trend (solid line) in the ex-vessel price spread index *b* (open circles) for reported May-August catches (inshore and offshore combined) of brown shrimp from Pensacola to the Mississippi River (statistical areas 10–12 combined), 1960–1981 (data from Tables 4 and 5, and Caillouet and Koi 1981). The 95% confidence bands are shown for the trend line by short dashes and for individual values of *b* by long dashes.

Figure 14. Trend (solid line) in the ex-vessel price spread index *b* (open circles) for reported May-August catches (inshore and offshore combined) of white shrimp from Pensacola to the Mississippi River (statistical areas 10–12 combined), 1960–1981 (data from Tables 4 and 5, and Caillouet and Koi 1981). The 95% confidence bands are shown for the trend line by short dashes and for individual values of *b* by long dashes.

TABLE 6.

Parameter estimates<sup>a</sup> of linear regressions of transformed cumulative weight (pounds, heads off) of catch,  $\ln P$ , on count, *C*, for reported May-August catches (inshore and offshore combined) of brown shrimp and white shrimp from the Texas coast (statistical areas 18–21 combined), 1979–1981.

Year	Brown Shrimp <sup>b</sup>			White Shrimp <sup>b</sup>		
	<i>c</i>	<i>d</i>	<i>r</i> <sup>2</sup>	<i>c</i>	<i>d</i>	<i>r</i> <sup>2</sup>
1979	20,674,053	-0.02112	0.973	3,332,356	-0.02300	0.890
1980	30,730,002	-0.02805	0.966	3,766,437	-0.04225	0.960
1981	62,640,964	-0.03768	0.962	2,577,937	-0.03633	0.983

<sup>a</sup>  $\ln(c)$  = intercept, *d* = slope, and *r*<sup>2</sup> = coefficient of determination.  
<sup>b</sup> Results of similar analyses of data from 1960 to 1978 can be found in Caillouet and Koi (1981).

TABLE 7.

Parameter estimates<sup>a</sup> of linear regressions of transformed cumulative weight (pounds, heads off) of catch,  $\ln P$ , on count, *C*, for reported May-August catches (inshore and offshore combined) of brown shrimp and white shrimp from the Mississippi River to Texas (statistical areas 13–17 combined), 1979–1981.

Year	Brown Shrimp <sup>b</sup>			White Shrimp <sup>b</sup>		
	<i>c</i>	<i>d</i>	<i>r</i> <sup>2</sup>	<i>c</i>	<i>d</i>	<i>r</i> <sup>2</sup>
1979	35,340,494	-0.00930	0.975	8,456,112	-0.04473	0.994
1980	21,930,962	-0.00516	0.962	6,638,058	-0.02483	0.977
1981	45,344,072	-0.00834	0.971	10,709,053	-0.03012	0.979

<sup>a</sup>  $\ln(c)$  = intercept, *d* = slope, and *r*<sup>2</sup> = coefficient of determination.  
<sup>b</sup> Results of similar analyses of data from 1960 to 1978 can be found in Caillouet and Koi (1981).

the Mississippi River, and highest for the Mississippi River to Texas (Tables 6–8; Figures 16, 18, and 20; see also Caillouet and Koi 1981). This indicated that the brown shrimp in the May-August catches from the Texas coast were consistently larger than those from Pensacola to the Mississippi River, and those from Pensacola to the Mississippi River were consistently larger than those from the Mississippi River to Texas. No such obvious differences in size of white shrimp in the May-August catches were apparent among the three coastal areas (Figures 17, 19, and 21).

To determine if there were significant trends in the index of size composition of the May-August catches, we calculated the linear regression of *d* on the last two digits (60–81) of each year, from 1960 to 1981, for both species and for each of the three coastal areas (Table 5). There were

significant upward trends in *d* for brown shrimp in all three coastal areas (Figures 16, 18, and 20), and for white shrimp from the Mississippi River to Texas (Figure 19), but there were no significant trends in *d* for white shrimp from the Texas coast (Figure 17) or for white shrimp from Pensacola to the Mississippi River (Figure 21). The significant upward trends for brown shrimp indicated that the size of brown shrimp in the reported May-August catches generally decreased in all three coastal areas from 1960 to 1981. The same was true for May-August white shrimp catches from the Mississippi River to Texas. The absence of significant trends in *d* for white shrimp catches from the Texas coast and from Pensacola to the Mississippi River indicated that there were no significant trends in size composition of the May-August catches in these two coastal areas. In these two

TABLE 8.

Parameter estimates<sup>a</sup> of linear regressions of transformed cumulative weight (pounds, heads off) of catch,  $\ln P$ , on count,  $C$ , for reported May-August catches (inshore and offshore combined) of brown shrimp and white shrimp from Pensacola to the Mississippi River (statistical areas 10–12 combined), 1979–1981.

Year	Brown Shrimp <sup>b</sup>			White Shrimp <sup>b</sup>		
	c	d	r <sup>2</sup>	c	d	r <sup>2</sup>
1979	13,586,810	-0.01427	0.849	707,215	-0.03047	0.936
1980	11,481,757	-0.02219	0.956	224,118	0.02600	0.926
1981	14,635,344	-0.01490	0.879	369,793	-0.03877	0.969

<sup>a</sup>  $\ln(c)$  = intercept,  $d$  = slope, and  $r^2$  = coefficient of determination.

<sup>b</sup> Results of similar analyses of data from 1960 to 1978 can be found in Caillouet and Koi (1981).




Figure 15. Relationship between the cumulative weight (millions of pounds, heads off) of catch,  $P$ , and count,  $C$ , for the reported May-August catch (inshore and offshore combined) of brown shrimp from the Texas coast (statistical areas 18–21 combined) in 1981 (graph A). In graph B, the line was fitted by linear regression (see Table 6).

cases, the mean of the indices from 1960 to 1981 was calculated as a horizontal line through the points (Figures 17 and 21). The data point for 1972 was excluded from calculation of the mean index for white shrimp from Pensacola to the Mississippi River (Figure 21), as in the previous section con-




Figure 16. Trend (solid line) in the size composition index  $d$  (open circles) for reported May-August catches (inshore and offshore combined) of brown shrimp from the Texas coast (statistical areas 18–21 combined), 1960–1981 (data from Tables 5 and 6, and Caillouet and Koi 1981). The 95% confidence bands are shown for the trend line by short dashes and for individual values of  $d$  by long dashes.




Figure 17. Mean (solid line) size composition index  $d$  (open circles) for reported May-August catches (inshore and offshore combined) of white shrimp from the Texas coast (statistical areas 18–21 combined), 1960–1981 (data from Table 6, and Caillouet and Koi 1981). The 95% confidence limits are shown for the mean by short dashes and for individual values of  $d$  by long dashes.

cerning the trend in price spread for white shrimp from this coastal area.

Within each coastal area, the size composition index  $d$  for brown shrimp catches in May-August 1981 fell within the 95% confidence bands for individual levels of  $d$  (Figures 16, 18, and 20), indicating that there was no significant departure in 1981 from the trend of decrease in size. In fact, though the size composition index for 1981 fell outside the 95% confidence interval for the trend lines (Figures 16, 18, and 20), it barely did so, emphasizing its closeness to the




Figure 18. Trend (solid line) in the size composition index *d* (open circles) for reported May-August catches (inshore and offshore combined) of brown shrimp from the Mississippi River to Texas (statistical areas 13-17 combined), 1960-1981 (data from Tables 5 and 7, and Caillouet and Koi 1981). The 95% confidence bands are shown for the trend line by short dashes and for individual values of *d* by long dashes.




Figure 19. Trend (solid line) in the size composition index *d* (open circles) for reported May-August catches (inshore and offshore combined) of white shrimp from the Mississippi River to Texas (statistical areas 13-17 combined), 1960-1981 (data from Tables 5 and 7, and Caillouet and Koi 1981). The 95% confidence bands are shown for the trend line by short dashes and for individual values of *d* by long dashes.




Figure 20. Trend (solid line) in the size composition index *d* (open circles) for reported May-August catches (inshore and offshore combined) of brown shrimp from Pensacola to the Mississippi River (statistical areas 10-12 combined), 1960-1981 (data from Tables 5 and 8, and Caillouet and Koi 1981). The 95% confidence bands are shown for the trend line by short dashes and for individual values of *d* by long dashes.




Figure 21. Mean (solid line) size composition index *d* (open circles) for reported May-August catches (inshore and offshore combined) of white shrimp from Pensacola to the Mississippi River (statistical areas 10-12 combined), 1960-1981 (data from Table 8, and Caillouet and Koi 1981). The 95% confidence limits are shown for the mean by short dashes and for individual values of *d* by long dashes.

continued trend of decreasing size. From 1979 to 1981, the index moved in a direction indicating increasing size of brown shrimp, but such variation was well within expected levels of variability in the index. The size composition index for white shrimp catches from the Mississippi River to Texas in May-August 1981 showed no significant departure from the trend of decrease in size (Figure 19). The size composition index for white

shrimp catches from the other two coastal areas in May-August 1981 fell within the 95% confidence limits for individual levels of *d* (Figures 17 and 21), showing that there was no significant departure in 1981 from the expected size composition. Thus the May-August 1981 size composition index was not a significant outlier in the 1960 to 1981 time series for either species in any of the three coastal areas.



*May-August cumulative ex-vessel value of catch by size category*

Caillouet and Koi (1981) used the logarithmic form of the following model to estimate parameters  $g$  and  $h$  by linear regression for each species, coastal area, and year from 1960 to 1978, and we updated this through 1981 (Tables 9–11):

$$\hat{D}_i = g(\exp hC_i) \quad (3)$$

where  $D_i$  = cumulative ex-vessel value (dollars) of catch in the  $i$ th size category. The ex-vessel values of the catches in each size category were cumulated starting with the size category of smallest shrimp and continuing toward the size category of largest shrimp, as was the case for cumulative pounds by size category, as described in the previous section. An example of the relationship between  $D$  and  $C$  is shown in Figure 22A, for brown shrimp from the Texas coast in 1981. An example of the fitted logarithmic form of model 3 is shown in Figure 22B. Very good fits were indicated by the very high coefficients of determination for each year (Tables 9–11, see also Caillouet and Koi 1981). All slopes,  $h$ , were negative, reflecting the construction of model 3 by cumulating ex-vessel value of catch from small-shrimp to large-shrimp size categories (Caillouet and Koi 1981). The slope,  $h$ , of each straight line is a simple index of the ex-vessel value composition of the May-August catch.

To determine whether there were significant trends in ex-vessel value composition of the May-August catch, we calculated the linear regression of  $h$  on the last two digits of each year from 1960 to 1981, for both species, and for each of the three coastal areas (Table 5). Only the upward trends in  $h$  for brown shrimp from the Texas coast (Figure 23) and for brown shrimp from Pensacola to the Mississippi River (Figure 27) were significant (Table 5). These upward trends indicated that ever increasing proportions of the ex-vessel value of the May-August catch of brown shrimp were represented by the size categories of smaller shrimp from these two coastal areas. The lack of significant trends in  $h$  for brown shrimp from the Mississippi River to Texas (Figure 25)

TABLE 9.

Parameter estimates<sup>a</sup> of the linear regressions of transformed cumulative ex-vessel value (dollars) of catch,  $\ln D$ , on count,  $C$ , for reported May-August catches (inshore and offshore combined) of brown shrimp and white shrimp from the Texas coast (statistical areas 18–21 combined), 1979–1981.

Year	Brown Shrimp <sup>b</sup>			White Shrimp <sup>b</sup>		
	$g$	$h$	$r^2$	$g$	$h$	$r^2$
1979	80,877,806	-0.03648	0.981	10,839,636	-0.03306	0.940
1980	109,104,860	-0.04202	0.978	13,595,100	-0.05468	0.975
1981	195,936,268	-0.05388	0.969	8,313,905	-0.05042	0.990

<sup>a</sup>  $\ln(g)$  = intercept,  $h$  = slope, and  $r^2$  = coefficient of determination.

<sup>b</sup> Results of similar analyses of data from 1960 to 1978 can be found in Caillouet and Koi (1981).

TABLE 10.

Parameter estimates<sup>a</sup> of linear regressions of transformed cumulative ex-vessel value (dollars) of catch,  $\ln D$ , on count,  $C$ , for reported May-August catches (inshore and offshore combined) of brown shrimp and white shrimp from the Mississippi River to Texas (statistical areas 13–17 combined), 1979–1981.

Year	Brown Shrimp <sup>b</sup>			White Shrimp <sup>b</sup>		
	$g$	$h$	$r^2$	$g$	$h$	$r^2$
1979	91,927,428	-0.01820	0.994	47,213,480	-0.06460	0.997
1980	40,696,392	0.00972	0.984	23,881,389	-0.04026	0.973
1981	78,311,890	-0.01689	0.980	43,460,368	-0.05200	0.960

<sup>a</sup>  $\ln(g)$  = intercept,  $h$  = slope, and  $r^2$  = coefficient of determination.

<sup>b</sup> Results of similar analyses of data from 1960 to 1978 can be found in Caillouet and Koi (1981).

TABLE 11.

Parameter estimates<sup>a</sup> of linear regressions of transformed cumulative ex-vessel value (dollars) of catch,  $\ln D$ , on count,  $C$ , for reported May-August catches (inshore and offshore combined) of brown shrimp and white shrimp from Pensacola to the Mississippi River (statistical areas 10–12 combined), 1979–1981.

Year	Brown Shrimp <sup>b</sup>			White Shrimp <sup>b</sup>		
	$g$	$h$	$r^2$	$g$	$h$	$r^2$
1979	41,494,855	0.02142	0.874	3,000,562	-0.04246	0.958
1980	35,357,108	-0.03169	0.967	809,434	-0.03933	0.921
1981	32,078,930	-0.02267	0.899	1,481,447	-0.05701	0.938

<sup>a</sup>  $\ln(g)$  = intercept,  $h$  = slope, and  $r^2$  = coefficient of determination.

<sup>b</sup> Results of similar analyses of data from 1960 to 1978 can be found in Caillouet and Koi (1981).

and for white shrimp from all three coastal areas (Figures 24, 26, and 28) indicated that the ex-vessel value composition of these catches exhibited no significant trend. In these cases, the mean of the indices from 1960 to 1981 was calculated as a horizontal line through the points (Figures 24, 25, 26, and 28). No data point for 1972 was included in calculation of the mean  $h$  for white shrimp from Pensacola to the Mississippi River (Figure 28), as in the previous section concerning size composition.

The levels of  $h$  for brown shrimp and white shrimp catches from all three coastal areas in May-August 1981 fell within the 95% confidence bands or confidence limits for individual levels of  $h$  (Figures 23–28), indicating that there was no significant departure in 1981 from the expected ex-vessel value composition. Thus, the ex-vessel value composition indices for May-August 1981 were not significant outliers in the 1960–1981 time series.

VALUE AND SIZE OF SHRIMP AS RELATED TO TEXAS CLOSURE

TABLE 12.

Simple correlation coefficients,  $r$ , for the relationships among indices<sup>a</sup> b, d and h and catch<sup>b</sup> for May-August catches (inshore and offshore combined) of brown shrimp and white shrimp from the Texas Coast (statistical areas 18-21 combined), the Mississippi River to Texas (statistical areas 13-17 combined), and Pensacola to the Mississippi River (statistical areas 10-12 combined), 1960-1981 (indices b, d and h are defined in the text).

Coastal Area/Species	Texas Coast						Mississippi River to Texas						Pensacola to the Mississippi River														
	Brown Shrimp			White Shrimp			Brown Shrimp			White Shrimp			Brown Shrimp			White Shrimp <sup>c</sup>											
	b	d	h	catch	b	d	h	catch	b	d	h	catch	b	d	h	catch	b	d	h	catch							
Texas Coast																											
Brown Shrimp	b	1.000	-0.314	-0.187	-0.157	0.604 <sup>d</sup>	0.123	0.221	-0.479 <sup>d</sup>	0.887 <sup>d</sup>	-0.233	0.230	0.230	-0.565 <sup>d</sup>	0.753 <sup>d</sup>	-0.186	0.065	-0.450 <sup>d</sup>	0.868 <sup>d</sup>	-0.333	-0.169	-0.498 <sup>d</sup>	0.777 <sup>d</sup>	0.249	0.349	0.447 <sup>d</sup>	0.052
Brown Shrimp	d	1.000	0.987 <sup>o</sup>	0.008	-0.095	0.421	0.355	0.182	-0.347	0.507 <sup>d</sup>	0.295	0.295	0.295	0.474 <sup>d</sup>	-0.227	0.096	-0.067	0.184	-0.456 <sup>d</sup>	0.542 <sup>d</sup>	0.441 <sup>d</sup>	-0.062	-0.201	-0.332	-0.462 <sup>d</sup>	-0.406	
Brown Shrimp	h	1.000	-0.007	-0.028	0.440 <sup>d</sup>	0.389	0.117	-0.253	0.5 <sup>d</sup>	0.360	0.360	0.360	0.430 <sup>d</sup>	-0.164	0.127	-0.011	0.115	0.374	0.528 <sup>d</sup>	0.438 <sup>d</sup>	-0.126	-0.133	-0.295	-0.415	-0.417	0.119	
Brown Shrimp	catch	1.000	1.000	0.285	0.424 <sup>d</sup>	-0.523 <sup>o</sup>	0.593 <sup>d</sup>	-0.168	0.072	-0.533	0.704 <sup>d</sup>	-0.217	-0.041	-0.302	0.563 <sup>d</sup>	-0.168	-0.083	-0.294	0.690 <sup>d</sup>	0.500 <sup>d</sup>	0.516 <sup>d</sup>	0.890 <sup>d</sup>	0.107	0.080	0.141	0.080	
White Shrimp	b	1.000	0.980 <sup>d</sup>	-0.143	0.249	-0.063	0.066	0.319	0.247	0.030	0.118	-0.107	0.219	0.029	0.087	0.093	0.093	0.256	0.090	0.129	-0.252	0.129	-0.252	0.129	-0.252	0.129	
White Shrimp	d	1.000	1.000	-0.270	0.315	-0.110	0.042	0.291	0.227	0.030	0.118	-0.107	0.219	0.029	0.087	0.093	0.093	0.256	0.090	0.129	-0.252	0.129	-0.252	0.129	-0.252	0.129	
White Shrimp	h	1.000	1.000	-0.402	0.252	0.143	0.660	0.247	0.030	0.118	-0.107	0.219	0.029	0.087	0.093	0.093	0.256	0.090	0.129	-0.252	0.129	-0.252	0.129	-0.252	0.129	-0.252	
White Shrimp	catch	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
Mississippi River to Texas																											
Brown Shrimp	b	1.000	-0.331	0.188	-0.587 <sup>d</sup>	0.866 <sup>d</sup>	-0.351	-0.049	-0.498 <sup>d</sup>	0.859 <sup>d</sup>	-0.435 <sup>d</sup>	0.726	-0.457 <sup>d</sup>	0.855 <sup>d</sup>	0.050	0.350	0.350	0.350	0.350	0.350	0.350	0.350	0.350	0.350	0.350	0.350	0.350
Brown Shrimp	d	1.000	0.834 <sup>d</sup>	0.176	1.000	-0.161	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
Brown Shrimp	h	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
Brown Shrimp	catch	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	
Pensacola to the Mississippi River																											
Brown Shrimp	b	1.000	-0.365	-0.442 <sup>d</sup>	0.809 <sup>d</sup>	0.304	0.563 <sup>d</sup>	0.142	0.169	-0.239	-0.155	-0.282	-0.196	1.000	0.965 <sup>d</sup>	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
Brown Shrimp	d	1.000	0.965 <sup>d</sup>	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
Brown Shrimp	h	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
Brown Shrimp	catch	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
White Shrimp <sup>c</sup>	b	1.000	-0.078	-0.068	0.991	-0.078	-0.068	0.991	-0.078	-0.068	0.991	-0.078	-0.068	0.991	-0.078	-0.068	0.991	-0.078	-0.068	0.991	-0.078	-0.068	0.991	-0.078	-0.068	0.991	-0.078
White Shrimp <sup>c</sup>	d	1.000	0.940 <sup>d</sup>	0.029	1.000	0.029	1.000	0.029	1.000	0.029	1.000	0.029	1.000	0.029	1.000	0.029	1.000	0.029	1.000	0.029	1.000	0.029	1.000	0.029	1.000	0.029	1.000
White Shrimp <sup>c</sup>	h	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000
White Shrimp <sup>c</sup>	catch	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000	1.000

a. The indices b, d, and h are shown in Tables 2-4, 6-8, and 9-11, respectively, for the years 1979-1981, and in Callouet and Kai (1981) for the years 1960-1978.

b. Total May-August catch, including "pieces," from Figures 2-7.

c. Data for 1972 were excluded from all correlation analyses involving indices b, d, and h for white shrimp from Pensacola to the Mississippi River (see Callouet and Kai 1981).

d. The correlation coefficient was significantly different from zero at the 95% level of confidence.




Figure 22. Relationship between the cumulative ex-vessel value (millions of dollars) of catch, D, and count, C, for reported May-August catch (inshore and offshore combined) of brown shrimp from the Texas coast (statistical areas 18-21 combined) in 1981 (graph A). In graph B, the line was fitted by linear regression (see Table 9).




Figure 23. Trend (solid line) in the ex-vessel value composition index  $h$  (open circles) for reported May-August catches (inshore and offshore combined) of brown shrimp from the Texas coast (statistical areas 18-21 combined), 1960-1981 (data from Tables 5 and 9, and Caillouet and Koi 1981). The 95% confidence bands are shown for the trend line by short dashes and for individual values of  $h$  by long dashes.




Figure 24. Mean (solid line) ex-vessel value composition index  $h$  (open circles) for reported May-August catches (inshore and offshore combined) of white shrimp from the Texas coast (statistical areas 18-21 combined), 1960-1981 (data from Table 9, and Caillouet and Koi 1981). The 95% confidence limits are shown for the mean by short dashes and for individual values of  $h$  by long dashes.




Figure 25. Mean (solid line) ex-vessel value composition index  $h$  (open circles) for reported May-August catches (inshore and offshore combined) of brown shrimp from the Mississippi River to Texas (statistical areas 13-17 combined), 1960-1981 (data from Table 10, and Caillouet and Koi 1981). The 95% confidence limits are shown for the mean by short dashes and for individual values of  $h$  by long dashes.

*Tests for correlations among indices b, d, and h, and the weight of the May-August catches*

In order to determine if indices  $b$ ,  $d$ , and  $h$  (Tables 2-4, 6-8 and 9-11, respectively; and Caillouet and Koi 1981) were closely related to one another, and if any of these indices were closely related to the total weight of the May-August catch (Figures 2-7), we calculated the simple correlation coefficients,  $r$ , for the relationships between all possible pairs of indices  $b$ ,  $d$ , and  $h$  and catch, for both




Figure 26. Mean (solid line) ex-vessel value composition index  $h$  (open circles) for reported May-August catches (inshore and offshore combined) of white shrimp from the Mississippi River to Texas (statistical areas 13-17 combined), 1960-1981 (data from Table 10, and Caillouet and Koi 1981). The 95% confidence limits are shown for the mean by short dashes and for individual values of  $h$  by long dashes.




Figure 27. Trend (solid line) in the ex-vessel value composition index  $h$  (open circles) for reported May-August catches (inshore and offshore combined) of brown shrimp from Pensacola to the Mississippi River (statistical areas 10-12 combined), 1960-1981 (data from Tables 5 and 11, and Caillouet and Koi 1981). The 95% confidence bands are shown for the trend line by short dashes and for individual values of  $h$  by long dashes.

species and each of the three coastal areas (Table 12). Each year provided one observation for each index and weight of catch. In the case of all correlation analyses involving indices  $b$ ,  $d$ , and  $h$  for white shrimp from Pensacola to the Mississippi River, one less observation was available, because the data for 1972 had to be excluded (Caillouet and Koi 1981).

The reader is cautioned that significant correlations (those significantly greater or less than zero) do not necessarily reflect cause and effect relationships. However, a




Figure 28. Mean (solid line) ex-vessel value composition index  $h$  (open circles) for reported May-August catches (inshore and offshore combined) of white shrimp from Pensacola to the Mississippi River (statistical areas 10-12 combined), 1960-1981 (data from Table 11, and Caillouet and Koi 1981). The 95% confidence limits are shown for the mean by short dashes and for individual values of  $h$  by long dashes.

number of general patterns emerged among the significant correlations (Table 12). For example, there were significant positive correlations between the May-August brown shrimp catches in all three coastal areas. Such results suggest that there was a region-wide influence on the May-August catches of brown shrimp, because there was a tendency for the catches to vary in the same direction from year to year in the three coastal areas. This could be explained by region-wide influences on brown shrimp recruitment, or by parallel changes in fishing effort in each of the three coastal areas, or both. The May-August white shrimp catches were not significantly correlated among the coastal areas, with the one exception: the white shrimp catches from the Mississippi River to Texas and from Pensacola to the Mississippi River were significantly correlated with each other. We offer no explanation for this correlation.

There were significant positive correlations between the ex-vessel price spread indices,  $b$ , for brown shrimp in all three coastal areas (Table 12). The same was true for price spread indices for white shrimp in all three coastal areas. Also, there were significant positive correlations between the price spread indices for the two species. This suggests a region-wide influence on price spread, because there was a tendency for the price spread to vary in the same direction from year to year in both species and in all three coastal areas. In most cases, but not all, there were significant negative correlations between the price spread index and the weight of the May-August catches (i.e., the current local supply of shrimp), both between species and between coastal areas, suggesting that price spread was wider when the catch was high, and narrower when the catch was low. Again, a regional influence was apparent, because the correlations existed between species and between coastal areas in most cases.

Among the highest of the significant correlations were the positive correlations between indices of size composition, *d*, and ex-vessel value composition, *h*, for a given species within a given coastal area. This was to be expected, because the ex-vessel value composition of the catch reflects, to a large extent, the size composition of the catch. This is particularly important, because it reinforces the general consensus that the size composition of the catch plays a major role in determining the ex-vessel value of the catch (Caillouet and Patella 1978).

While there were a number of additional significant correlations depicted in Table 12, they did not follow any particular pattern; therefore we did not attempt an explanation for these cases. However, the absence of significant correlations is noteworthy in at least one case. For example, within a given species and coastal area, there were no significant correlations between the weight of the May-August catch and the size composition index. This indicates that the size composition of the catch is not the major factor affecting the weight of the catch, or vice versa. Year to year variations in recruitment, fishing effort or both, may be overriding factors in this regard. That is not to say that size composition has no effect on the weight of the catch. As might be expected, differences in fishing regulations and fishing strategy among the three coastal areas have had pronounced effects upon the size composition of the catches (Caillouet et al. 1980, Caillouet and Koi 1980).

#### DISCUSSION

Based upon our analyses of the best catch statistics available from the TIMS in February 1982, we detected significant trends of decrease in size of brown shrimp in the reported May-August catches from the three coastal areas over the years 1960 to 1981. This corroborates similar findings by Caillouet and Koi (1980) for the period 1960–1978. In addition, Caillouet et al. (1980) detected significant decreases in size of brown shrimp in reported annual catches from Texas and Louisiana from 1959 to 1976, and Caillouet and Koi (1980) detected significant decreases in size of brown shrimp in reported annual landings from the northern Gulf from 1961 to 1977.

In 1981, the size composition index for the May-August brown shrimp catch in each of the three coastal areas did not depart significantly from that expected, based on the general trends of decrease in size from 1960 to 1981 (Figures 16, 18, and 20). It barely fell out of the 95% confidence bands for the trend lines, and it was well within the 95% confidence intervals for the individual levels of the index predicted for each coastal area for 1981, based on the entire 1960 to 1981 time series. Despite these trends of decrease in size, the total weight of the May-August catch of brown shrimp from all three coastal areas was higher in 1981 than in 1980 (Figures 2, 4, and 6), and the ex-vessel value of the May-August catch of brown shrimp from the Texas coast

was at an all time high (Figure 2). The higher catches of brown shrimp in 1981, coupled with the continued increase in ex-vessel price spread, enhanced the ex-vessel value of the brown shrimp catches, particularly in Texas where brown shrimp continued to be larger than those in catches from the Mississippi River to Texas and Pensacola to the Mississippi River.

The May-August brown shrimp catches in all three coastal areas showed a tendency to vary in the same direction from year to year. Also, the lack of a significant correlation, using the 1960 to 1981 time series, between the total weight and the size composition of the May-August catch of brown shrimp within each of the three coastal areas, suggested that size composition was not the major determinant of weight of the catch or vice versa. These findings, coupled with the fact that there were no detectable departures in 1981 from the trends of decreasing size in brown shrimp in the May-August catches from the three coastal areas during 1960 to 1981, suggested that the observed increases in brown shrimp catch from all three coastal areas reflected some region-wide influence.

According to Jones and Zweifel (1982), there was more fleet movement in June-August 1981 than in a comparable period in 1980, and they attributed this to the Texas Closure. The change in mobility included shifts to areas away from the Texas coast during the closure and back to the Texas coast thereafter. Klima et al. (1982) indicated that recruitment from Texas bays to the offshore fishery appeared to be average to good in 1981, but not so good as to account for the outstanding abundance levels found offshore of Texas. According to Charles J. White (Louisiana Department of Wildlife and Fisheries, Lake Charles, Louisiana, personal communication, March 1982), brown shrimp recruitment in Louisiana, both west and east of the Mississippi River, was near record levels in 1981. Also, J. Y. Christmas (Gulf Coast Research Laboratory, Ocean Springs, Mississippi, personal communication, March 1982) indicated that there was good brown shrimp recruitment in 1981 in the area of Mississippi Sound off the coast of Mississippi.

There were no significant trends in size composition of reported May-August catches of white shrimp from the Texas coast and from Pensacola to the Mississippi River from 1960 to 1981, a result similar to that reported for May-August catches from 1960 to 1978 by Caillouet and Koi (1981). However, there was a significant trend of decrease in size of white shrimp in the May-August catch from the Mississippi River to Texas from 1960 to 1981. The size composition of the May-August 1981 catch of white shrimp in the three coastal areas showed no significant departure from that expected based on 1960–1981 trends or mean levels (Figures 17, 19, and 21). On the other hand, Caillouet et al. (1980) and Caillouet and Koi (1980) detected decreases in size of white shrimp in reported annual catches and landings, respectively, from the northern Gulf of Mexico.

Therefore, the observed trends of decrease in size of white shrimp in the annual catches and landings from the northern Gulf must reflect an overriding influence of catches in months other than May-August. The low percentages of the annual white shrimp catches taken in May-August (Table 1) corroborate this. While the ex-vessel price spread for white shrimp in the May-August catch from the Mississippi River to Texas and from Pensacola to the Mississippi River increased from 1960–1981, no such trend was detected in price spread for white shrimp from the Texas coast (Table 5).

Our analyses did not demonstrate immediate and detectable increases in size of brown shrimp or white shrimp in the May-August catch concomitant with the Texas Closure in 1981. However, the total weight and ex-vessel value of the May-August catches of both species in each coastal area were higher in 1981 than in 1980, with one exception: white shrimp from the Texas coast exhibited a slight decrease in weight of May-August catch in 1981 as compared to 1980, but there was nevertheless an increase in ex-vessel value of this catch.

The magnitude of the increase in weight of the May-August catch of brown shrimp between 1980 and 1981 was greater for the Mississippi River to Texas (18.6 million pounds, or 94%, Figure 4) than for either the Texas coast (12.1 million pounds, or 69%, Figure 2) or Pensacola to Mississippi River (3.2 million pounds, or 45%, Figure 6). This occurred despite the fact that approximately half of statistical area 17, which is located within the Mississippi River to the Texas coastal area, was closed to shrimping during the Texas Closure. Also, the magnitude of the increase in ex-vessel value of the May-August catch of brown shrimp between 1980 and 1981 was greater for the Mississippi River to Texas (22.6 million dollars, or 66%, Figure 4) than for either the Texas coast (18.2 million dollars, or 38%, Figure 2) or Pensacola to the Mississippi River (0.9 million dollars, or 5%, Figure 6). Nonetheless, in May-August 1981, the Texas coast continued to produce brown shrimp catches of higher ex-vessel value than the other two coastal areas, because these catches were represented by larger shrimp which brought a higher price than in the other two coastal areas. Our findings are consistent with the findings of Poffenberger (1982) who estimated an increase in revenue of 21.5 million dollars to the Gulf of Mexico brown shrimp fishery during May-August 1981.

We concluded that the increases in brown shrimp catches in May-August 1981, as compared to May-August 1980, appeared to be due to a region-wide influence, perhaps good recruitment in 1981, or increased fishing effort, or both. This finding is not necessarily inconsistent with those of Klima et al. (1982), Nichols (1982), or Matthews (1982),

because these authors dealt only with offshore catches, and concluded that offshore abundance and yield of brown shrimp was enhanced by the closure. Our analyses dealt with inshore and offshore catches combined. It was obvious that the continued increase in ex-vessel price spread among size categories of shrimp continued to enhance the ex-vessel value of the catches, especially on the Texas coast.

The reader is reminded that our analyses dealt only with the May-August catches. For the Texas coast, the May-August period produces a smaller proportion (averaging 58%) of the calendar year annual brown shrimp catch from the Texas coast, than does a similar period for the Mississippi River to Texas (81%), or for Pensacola to the Mississippi River (83%). Thus, the May-August time interval appears to be too brief a period to show the full effects of the Texas Closure on size composition or yield of the brown shrimp. An examination of the annual catches, fishing effort, and recruitment in 1981, as compared to earlier years, might be useful in more fully assessing the impacts of the Texas Closure. An analysis applying our approach to biological year catch statistics for brown shrimp might also be useful in further assessments of impacts of the Texas Closure or other management strategies. For that purpose, we would recommend a biological year beginning in May of one year and ending in April of the following year, based upon our examination of the weight and size composition of monthly catches of brown shrimp.

#### ACKNOWLEDGMENTS

We are especially grateful to those who designed and implemented the gathering of data on weight and ex-vessel value of shrimp catches by statistical area, species and size category, since they have made our analyses possible. Notable among them are Charles H. Lyles, Director, Gulf States Marine Fisheries Commission (formerly of the USFWS); George W. Rounsefell (deceased; formerly Director, Galveston Biological Laboratory, BCF, USFWS); Joseph H. Kutkuhn, Director, Great Lakes Fisheries Laboratory, USFWS, Ann Arbor, Michigan (formerly Assistant Director, Galveston Biological Laboratory, BCF, USFWS); George W. Snow (retired; formerly Chief, Division of Statistics and Market News, NMFS, New Orleans, Louisiana); and TIMS.

The manuscript was reviewed by Dr. Edward F. Klima, Director, NMFS Southeast Fisheries Center (SEFC), Galveston Laboratory, and by Dr. Albert C. Jones, Dr. Joseph Powers and Mr. John R. Poffenberger, NMFS, SEFC, Miami, Florida, who provided many helpful suggestions. Beatrice Richardson, clerk-typist, NMFS, SEFC, Galveston Laboratory, typed the manuscript.

#### REFERENCES CITED

- Caillouet, C. W. & D. B. Koi. 1980. Trends in ex-vessel value and size composition of annual landings of brown, pink, and white shrimp from the Gulf and south Atlantic coasts of the United States. *Mar. Fish. Rev.* 42(12):18–27.
- \_\_\_\_\_. 1981. Trends in ex-vessel value and size composition of reported May-August catches of brown and white shrimp from